

University of California Immunization Requirements

Welcome New Graduate Division Students.

The University of California is committed to protecting the health and well-being of our students. Therefore, all UC campuses are implementing procedures to ensure that students are educated about and receive vaccinations to prevent potentially serious and contagious diseases. Despite the fact that many people receive the recommended vaccines, there are still documented outbreaks of vaccine-preventable-diseases (VPD) in California each year amongst those who were not completely immunized. Information about these VPDs is provided on the following page.

You are required to obtain and submit proof of immunity to these diseases and complete a TB screening questionnaire via the MyHealthRecord.ucsf.edu portal.

Vaccination and Screening Requirements and Recommendations are adopted from the California Department of Public Health IMMUNIZATION & SCREENING RECOMMENDATIONS FOR COLLEGE STUDENTS in place February 9, 2016; NOTE that revisions to the CDPH recommendations will be reflected in UC requirements for the subsequent academic term

All incoming UCSF students are REQUIRED to have received these vaccines and undergo Tuberculosis screening
<i>Required Vaccination or Screening</i>
Measles, Mumps and Rubella (MMR)
Varicella (chickenpox)
Tetanus, Diphtheria and Pertussis (Tdap)
Screening for Tuberculosis

Additional immunizations STRONGLY RECOMMENDED for preventive care include:

- Influenza vaccine (annual)
- Hepatitis A vaccine
- Hepatitis B vaccine
- Human papillomavirus vaccine (for women through age 26 years; males through age 26 years may also receive the quadrivalent HPV vaccine)
- Pneumococcal vaccine (for students with certain medical conditions e.g., severe asthma, diabetes, chronic liver or kidney disease)
- Poliovirus vaccine (if series not completed as a child)
- Meningococcal conjugate (for graduate students new to residential housing)
- Meningococcal B (for students aged ≥ 10 years who are at increased risk for meningococcal disease- e.g. those with persistent complement component deficiencies; persons with anatomic or functional asplenia; microbiologists routinely exposed to isolates of *Neisseria meningitidis*; and persons identified to be at increased risk because of a serogroup B meningococcal disease outbreak **OR** those ages 16-23 who elect vaccination after discussion with their healthcare provider)
- Vaccines for international travel, based on destination

University of California Immunization Requirements

Vaccine Preventable Diseases—What you should know about the diseases:

Many VPD's were very common before widespread vaccination programs began. With decreasing vaccination rates, outbreaks of some of these infections have started to recur.

Diphtheria	Causes a thick covering in the back of the throat. It can lead to breathing problems, paralysis, heart failure, and even death.
Hepatitis A	A serious liver disease that is usually spread by close personal contact or by eating contaminated food or water. Hepatitis A can cause mild, "flu-like" illness, jaundice (yellow skin or eyes), and severe stomach pains and diarrhea. Teens and adults are most likely to have a severe case.
Hepatitis B	Another serious liver disease that can cause short-term (acute) appetite loss, diarrhea, vomiting, fatigue, jaundice (yellow skin or eyes), and/or pain in muscles, joints, and stomach. It also can cause long-term (chronic) illness that leads to liver damage (cirrhosis), liver cancer, and death.
Human Papillomavirus	A virus that causes genital warts and a variety of cancers, and is spread by direct skin-to-skin contact.
Influenza (flu)	A contagious viral disease spread through coughing or sneezing. Influenza can lead to pneumonia, sinus or ear infections, worsening of chronic conditions such as asthma or diabetes, and even death.
Measles	This virus causes a rash, cough, runny nose, eye irritation, and fever. It can lead to ear infections, pneumonia, seizures, brain damage, and even death.
Meningococcal Disease	A typically severe bacterial infection that can cause hearing loss, learning problems, brain damage, or loss of limbs. About 1 in 10 people with meningococcal disease will die from it. The disease is spread when people live in close contact or through other close contact such as partying or kissing. Teens and young adults have a higher risk for meningococcal disease.
Mumps	This virus causes fever, headache, swollen glands and painful swelling of the testicles or ovaries. It can lead to deafness, meningitis (infection of the brain and spinal cord covering), and, rarely, death.
Pertussis	Also called <i>whooping cough</i> , causes prolonged coughing spells for weeks to months that can end in vomiting. It can lead to pneumonia, seizures (jerking and staring spells), brain damage, and death.
Polio	A disease caused by a virus which can cause paralysis (cannot move arms or legs). It can kill people who get it, usually by paralyzing the muscles that help them breathe.
Pneumococcal disease	A leading cause of bacterial meningitis in the United States. It causes many health problems in, including pneumonia, meningitis, blood infections, sinus infections, ear infections, and even death.
Rubella	Also known as <i>German Measles</i> , this virus usually causes mild fever and a rash. However, if a woman gets rubella while pregnant, she could have a miscarriage or her baby could have serious birth defects.
Tetanus	Also called <i>Lockjaw</i> , this is a painful tightening of the muscles, usually all over the body. It can lead to "locking" of the jaw so the person cannot open their mouth or swallow. Tetanus can be fatal.
Tuberculosis	A disease caused by a bacterium called <i>Mycobacterium tuberculosis</i> . The bacteria usually attack the lungs, but can infect any part of the body, such as the kidney, spine, and brain. If not treated properly over several months, tuberculosis can be fatal.
Varicella	Also known as <i>Chickenpox</i> , this virus causes a rash, itching, fever, and tiredness. While chickenpox is usually a mild illness, it also can lead to severe skin infections, scars, pneumonia, brain damage, or death.